



UNIVERSITY OF  
**FLORIDA**

Circular 1001

Cooperative Extension Service  
Institute of Food and Agricultural Sciences

## Managing Pesticides for Rice Production and Water Quality Protection<sup>1</sup>

A.G. Hornsby, T. M. Buttler, J.A. Dusky, F.J. Coale and L.E. Datnoff<sup>2</sup>

### WATER QUALITY CONSIDERATIONS IN RICE PRODUCTION

Concern about the harmful effects of pesticides on surface water and groundwater quality should motivate rice producers to select pesticides with the least potential to cause water quality problems. Many rice producers live in rural areas near where they and other producers grow rice, therefore, their personal water supply is susceptible to contamination. Unfortunately, information that allows producers to select pesticides less likely to affect water quality has not previously been readily available.

Our purpose is to provide information that can help producers select pesticides that will have a minimum adverse impact on water quality. The procedure considers the soil properties of the application site, the mobility and persistence of pesticides in these soils, and the toxicity of the pesticides in water to humans and aquatic species. A proper selection will decrease chances of adversely affecting surface water and groundwater quality. Certain combinations of soil and pesticide properties (along with weather conditions) can pose a significant potential hazard to water quality.

Our goal is to identify and avoid these circumstances. Information contained in this circular can help rice producers make better decisions about the pesticides that they use. This document in no way endorses any particular pest control product. All products must be used in accordance with the label.

### MATERIALS NEEDED TO USE THIS PROCEDURE

To effectively use this procedure you will need the following source materials:

- 1) A copy of the current IFAS Pest Management Guides or other appropriate information sources that identify pesticides that control specific pests.
- 2) A copy of your county soil survey report to identify the soil types found in your fields.
- 3) A copy of the Soil Science Fact Sheet entitled "[Name of your county]: Soil Ratings for Selecting Pesticides" for your county, available from your county Cooperative Extension Office. The basis of these ratings are given in the IFAS Extension

1. This document is Circular 1001, Soil and Water Science Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. First published: May 1991. Last revision: January 1998. Please visit the FAIRS Website at <http://hammock.ifas.ufl.edu>.
2. A.G. Hornsby and T.M. Buttler, Soil and Water Science Department; J.A. Dusky, F.J. Coale, and L.E. Datnoff, Everglades Research and Extension Center, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, 32611.

COOPERATIVE EXTENSION SERVICE, UNIVERSITY OF FLORIDA, INSTITUTE OF FOOD AND AGRICULTURAL SCIENCES, Christine Taylor Waddill, Director, in cooperation with the United States Department of Agriculture, publishes this information to further the purpose of the May 8 and June 30, 1914 Acts of Congress; and is authorized to provide research, educational information and other services only to individuals and institutions that function without regard to race, color, age, sex, handicap or national origin. The information in this publication is available in alternate formats. Single copies of extension publications (excluding 4-H and youth publications) are available free to Florida residents from county extension offices. Information on copies for out-of-state purchase is available from Publication Distribution Center, University of Florida, P. O. Box 110011, Gainesville, FL. 32611-0011. Information about alternate formats is available from Educational Media and Services, University of Florida, P. O. Box 110810, Gainesville, FL. 32611-0810. This information was published May 1991 as Cir 1001, Florida Cooperative Extension Service. Revised January 1998.

Circular 959 entitled "Soil Ratings for Selecting Pesticides for Water Quality Goals," which is also available from your county Cooperative Extension Office.

Note: If your county has not yet been mapped by the Natural Resources Conservation Service (formerly the Soil Conservation Service), you will need to contact the local NRCS office for a site evaluation and determination of soil types and ratings for leaching and runoff of pesticides.

## IMPORTANT FACTORS THAT AFFECT PESTICIDE SELECTION

How pesticides behave in the soil is determined by many factors including properties of the pesticides and of the soil at the application site. Some of the factors that should be considered when selecting pesticides with minimal potential for water quality impacts are:

### Pesticide Properties

- 1) The organic carbon adsorption coefficient,  $K_{oc}$ , describes the relative affinity or attraction of the pesticide to soil materials and therefore its mobility in the soil.
- 2) The biological degradation half-life,  $T_{1/2}$ , is a measure of persistence of the pesticide in soil.
- 3) The maximum contaminant level (MCL), or the lifetime health advisory level (HAL), or equivalent (HALEQ), is a measure of health risk to humans of pesticide contaminated drinking water.
- 4) Aquatic toxicity,  $LC_{50}$ , is a measure of the ability of the pesticide to cause 50% mortality in aquatic test species.

### Soil Properties

- 1) Hydraulic permeability is a measure of the soils ability to allow water to percolate through it.
- 2) Organic matter is important for providing binding sites for pesticides, thus reducing their mobility and increasing their opportunity to be degraded by soil microorganisms.

- 3) Slope affects the potential for water to run off the land surface.

### Management Practices

- 1) Pesticide application frequencies and rates determine the total amount applied. Lower frequencies and rates reduce the potential for contamination.
- 2) Application methods affect the amount of pesticide subject to transport by water. For example, if applied directly to the soil, there is a greater probability that more of the product will be available for leaching or runoff than if applied to the foliage. If the product is incorporated into the soil, leaching may be the most important loss pathway. Pesticides applied to the foliage may be lost to the atmosphere, decomposed by sunlight, or absorbed by the foliage, thereby reducing the amount available for wash-off and transport to water bodies.
- 3) Irrigation practices can also determine the loss pathways of pesticides. Pesticides often move with water, so the less excess water that is applied, the less potential there is for a pesticide to move past the crop root zone or to run off in surface water. Rainfall or overhead irrigation can wash off significant quantities of pesticides from foliage immediately after application.

## INDICES USED TO SELECT PESTICIDES

Table 1 contains two important indices, the pesticide leaching potential (RLPI) and the pesticide runoff potential (RRPI). Both indices are relative. For a given soil, these indices rank the pesticides by their potential to move from the application site by the indicated pathway (leaching or runoff). The indices are based on the organic carbon sorption coefficient and degradation half-life values of each pesticide. Values for these parameters have been taken from scientific literature, technical manuals, and company product literature.

### Relative Leaching Potential Index

The Relative Leaching Potential Index (RLPI) defines the relative attenuation (reduction in mass as it moves through the soil) of each pesticide in soil, and therefore its potential to leach to groundwater. Pesticides that are very mobile, for example, those that have  $K_{oc}$  values less than 100 in sandy soils, or 50 or less in fine-textured soils should be used with caution. There is some uncertainty in the data used to calculate this index. However, since the values are relative they can still be used. It is important to realize that the smaller the RLPI value of a pesticide, the greater is its potential to leach.

### Relative Runoff Potential Index

The Relative Runoff Potential Index (RRPI) defines the relative immobility and availability of each pesticide in soil, and therefore its potential to remain near the soil surface and be subject to loss in the aqueous phase or sediment phase of runoff. There is some uncertainty in the data used to calculate this index. However, since the values are relative they can still be used. The smaller the RRPI value of a pesticide, the greater is its potential to be lost in runoff.

### Maximum Contaminant Level, Lifetime Health Advisory Level or Equivalent

Table 1 also contains information on the toxicity of pesticides to humans and aquatic species. This information can be used as a secondary consideration in the pesticide selection procedure.

The Maximum Contaminant Level, MCL, is the highest allowable concentration in drinking water supplied by municipal water systems. It is a Primary Drinking Water Standard based on health considerations and is enforceable by the USEPA. Pesticides that may potentially cause chronic health effects such as cancer, birth effects, miscarriages, nervous system disorders or organ damage are assigned a MCL value by the USEPA. Although the MCL is usually expressed in milligrams per liter (mg/L), in this circular it will be expressed as micrograms per liter ( $\mu\text{g/L}$ , or ppb). The Lifetime Health Advisory Level, HAL, provides a measure of pesticide toxicity to humans. The HAL as defined by the USEPA is the concentration of a chemical in drinking water that is not

expected to cause any adverse health effects over a lifetime of exposure (70 years), with a margin of safety. If the chemical has not been assigned a MCL or HAL value, a health advisory level equivalent, HALEQ, (denoted by an asterisk) has been calculated using the same formula as the USEPA ( $\text{HALEQ} = R_f D \times 7000$ ) where  $R_f D$  is the reference dose determined by the USEPA. For non-carcinogenic pesticides the calculated HALEQ should not differ by more than a factor of 10 from the value forthcoming from the USEPA. The Hal and the HALEQ have units of micrograms per liter ( $\mu\text{g/L}$  or ppb). If a pesticide has a MCL value assigned, we use that value rather than the HAL or HALEQ. The smaller the value the greater is the toxicity to humans.

### Aquatic Toxicity

The Aquatic Toxicity provides a measure of pesticide toxicity to aquatic species. The values given in Table 1 are the lethal concentrations at which 50% of the test species die ( $LC_{50}$ ). Unless otherwise noted by a lower case letter following the value, the test species was rainbow trout. The smaller the value the greater is the toxicity to aquatic species.

Data for  $K_{oc}$ , RLPI, RRPI, MCL/HAL/HALEQ, and aquatic toxicity are given for the active ingredient (common name) of a product. When using a product that is a mixture of two or more active ingredients use the RLPI, RRPI, MCL/HAL/HALEQ, and Aquatic Toxicity value for the most restrictive active ingredient in the mixture.

**Important Note:** The information presented in Table 1 **DOES NOT** supersede or replace the information on the pesticide container label or product literature.

## PROCEDURE FOR SELECTING PESTICIDES TO REDUCE ADVERSE WATER QUALITY IMPACTS

A "Pesticide Selection Worksheet" is provided as a convenient way to organize the information needed to select pesticides to avoid water pollution by pesticides in a particular production or management unit. Instructions for using the worksheet are provided. The function of the worksheet is to match the soil leach and runoff ratings at the application site with the pesticide RLPI

(leaching) and RRPI (runoff) indices and toxicity values given in Table 1.

This will indicate the relative potential for pesticides to leach or run off from a particular site and consider the toxicity of the pesticides to humans or aquatic life if the pesticides leach into groundwater or if runoff enters surface impoundments or streams. The last two columns are for recording the rice producer's choices and reasons for selecting particular products.

Our intent is to provide a decision support tool for the rice producer. The producer is responsible for making the final choice. The completed worksheet can serve as a permanent record of the selection process used and decision made by the producer

### Using the Worksheet

- 1) **Target Pest:** Correct identification of the pests that need to be controlled is essential! Check with knowledgeable experts and utilize competent diagnostic laboratories so that a proper diagnosis can be made. Misdiagnosis results in the wasteful use of unnecessary pesticides and needless increases in production costs. List confirmed pests in column 1 of the PESTICIDE SELECTION WORKSHEET.
- 2) **Recommended Pesticides:** Use the current IFAS Pest Management Guides, or other appropriate information sources to identify the pesticides that control the pests of concern. List these pesticides in column 2 of the PESTICIDE SELECTION WORKSHEET.
- 3) **Pesticide Properties:** For each pesticide listed in column 2 on the PESTICIDE SELECTION WORKSHEET, copy the numeric value for  $K_{oc}$ , RLPI, RRPI, MCL/HAL/HALEQ, and Aquatic Toxicity from Table 1 into columns 3, 4, 5, 6, and 7 of the PESTICIDE SELECTION WORKSHEET.
- 4) **Soil Properties:** Consult the County Soil Survey Report soil map sheets to locate your production fields and to identify the soils that occur in these fields. Use the Soil Science Fact Sheet entitled "[Your County]: Soil Ratings for Selecting

Pesticides" (available from your county Cooperative Extension Office) to determine the leaching and surface runoff rating of the soils in your fields. As you determine the soil leach rating and the soil runoff rating for each soil in each field, list the soil name, soil leach rating, and soil runoff rating in columns 8, 9, and 10, respectively, of the PESTICIDE SELECTION WORKSHEET.

- 5) **Selection of Pesticides:** Using information that you have compiled on the PESTICIDE SELECTION WORKSHEET, select appropriate pesticides using the **Pesticide Selection Criteria** to match soil and pesticide properties. The selection made can be recorded in column 11 and notes relating to the selection can be recorded in column 12.

#### Notes:

- 1) If the pesticide product selected is a formulated mixture or a tank mix, each active ingredient must be considered. The most restrictive pesticide in the mixture will determine the choice. Trade names in Table 1 followed by (M) are formulated mixtures.
- 2) Sometimes there may not be a clear choice from among the alternative chemicals available to control a particular pest. In these cases, first order screening using the RLPI or RRPI only can suffice.
- 3) Depth to groundwater and local geohydrology may influence your final selection. Shallow groundwater is more vulnerable to contamination. Deep water tables with intervening impermeable geologic layers are much less vulnerable.
- 4) Distance to surface water bodies may also influence your final selection. Surface waters adjacent to or near the pesticide application site are more vulnerable to contamination than those further away. If surface runoff from the application site usually infiltrates into the soil off site before reaching a surface water body, then the MCL/HAL/HALEQ should be considered as the secondary screening index.

**Criteria for Matching Soil Ratings with Pesticide Indices**

Pesticides with less potential to adversely affect water quality can be selected by matching the soil ratings and pesticides with the following criteria:

PESTICIDE SELECTION CRITERIA				
IF SOIL RATINGS ARE:		THEN		
LEACH	RUNOFF	SELECT PESTICIDE WITH:		
HIGH	LOW	Larger RLPI value,	<u>AND</u>	Larger MCL/HAL/HALEQ value.
MEDIUM	LOW	Larger RLPI value,	<u>AND</u>	Larger MCL/HAL/HALEQ value.
LOW	LOW	Larger RLPI and RRPI values,	<u>AND</u>	Larger MCL/HAL/HALEQ and Aquatic Toxicity values.
HIGH	MEDIUM	Larger RLPI and RRPI values,	<u>AND</u>	Larger MCL/HAL/HALEQ and Aquatic Toxicity values.
MEDIUM	MEDIUM	Larger RLPI and RRPI values,	<u>AND</u>	Larger MCL/HAL/HALEQ and Aquatic Toxicity values.
LOW	MEDIUM	Larger RRPI value,	<u>AND</u>	Larger Aquatic Toxicity value.
HIGH	HIGH	Larger RLPI and RRPI values,	<u>AND</u>	Larger MCL/HAL/HALEQ and Aquatic Toxicity values.
MEDIUM	HIGH	Larger RRPI and RLPI values,	<u>AND</u>	Larger Aquatic Toxicity and MCL/HAL/HALEQ values.
LOW	HIGH	Larger RRPI value,	<u>AND</u>	Larger Aquatic Toxicity value.

**ACKNOWLEDGMENTS**

The development of this document was supported by the USDA/ES Water Quality Initiative Project #89EWQI-1-9134 and the Center for Natural Resources, University of Florida, Gainesville, FL.

**Table 1.** Rice-Pesticide parameters for selecting pesticides to minimize water quality problems.

Trade name <sup>1</sup>	Common name	Application Type <sup>2</sup>		Sorptions Coefficient <sup>3</sup>		Relative Losses		Toxicity		
		Soil	Foliar	K <sub>oc</sub> (ml/g)		Leaching RLPI <sup>4</sup>	Runoff RRPI <sup>5</sup>	MCL, HAL or HALEQ <sup>6</sup> (ppb)	Aquatic LC <sub>50</sub> <sup>7</sup> (ppm)	
<b>Herbicides</b>										
2,4-D amine	2,4-D dimethylamine salt		x	20	E	20	20	70		100
Basagran	bentazon		x	34		nd	nd	0		510
Blazer	acifluorfen		x	119		nd	nd	0		17
Bolero	thiobencarb	x		900		428	52	70	*	1.2
Prowl	pendimethalin	x		5000		555	2	300	*	0.199b
Roundup	glyphosate amine salt		x	24000	E	>2,000	1	700		8.3
Stam	propanil		x	149		1490	>1,000	40	*	6.13a
Weedone	MCPA ester		x	1000	E	400	40	11		232
<b>Insecticides/Miticides</b>										
Cythion	malathion		x	1800		>2,000	555	200		0.2
Sevin	carbaryl		x	300		300	300	700		114
<b>Fungicides</b>										
Apron	metalaxyl	x		50		7	7	400	*	>100
Benlate	benomyl		x	1900		283	7	350	*	0.17
Dithane	mancozeb	x		2000	E	285	7	20	*	1b
Manzate	mancozeb	x		2000	E	285	7	20	*	1b
Rovral	iprodione		x	700		500	102	300	*	6.7
Thiolux	sulfur		x	nd		nd	nd	nd		non toxic
Thiram	thiram	x		670		446	99	40	*	0.13

**Table 1.** Rice-Pesticide parameters for selecting pesticides to minimize water quality problems.

Trade name <sup>1</sup>	Common name	Application Type <sup>2</sup>		Sorption Coefficient <sup>3</sup>		Relative Losses		Toxicity		
		Soil	Foliar	K <sub>oc</sub> (ml/g)		Leaching RLPI <sup>4</sup>	Runoff RRPf <sup>5</sup>	MCL, HAL or HALEQ <sup>6</sup> (ppb)		Aquatic LC <sub>50</sub> <sup>7</sup> (ppm)
Tilt	propiconazole		x	650	E	59	13	100	*	1.3b
Vitavax	carboxin	x		260		866	866	700		2
<sup>1</sup> Tradename:		(M) indicates that the product is a mixture of two or more active ingredients.								
<sup>2</sup> Application Type:		INC: incorporated; INJ: injected; PRE: preemergence; X: applied to soil surface or foliage								
<sup>3</sup> Sorption Coefficient:		E: estimated G: educated guess								
<sup>4</sup> Relative Leaching Potential Index (RLPI):		Smaller number indicates greater leaching hazard.								
<sup>5</sup> Relative Runoff Potential Index (RRPI):		Smaller number indicates greater runoff hazard.								
<sup>6</sup> Drinking Water:		Maximum Contaminant Level (MCL), Lifetime Health Advisory Level (HAL); * Lifetime Health Advisory Level Equivalent (HALEQ);								
<sup>7</sup> Aquatic Toxicity LC <sub>50</sub> :		value is for rainbow trout 48 or 96 hr exposure time, unless otherwise specified. a=channel catfish b=bluegill c=carp j=fat head minnow								
nd: no data available										

PESTICIDE SELECTION WORKSHEET

Landowner/Operator Name: \_\_\_\_\_ County: \_\_\_\_\_ Date: \_\_\_\_\_

Crop: \_\_\_\_\_ Farm ID: \_\_\_\_\_ Field ID \_\_\_\_\_ Sheet \_\_\_\_\_ of \_\_\_\_\_

Target Pest (1)	IFAS Recommended Pesticides (2)	K <sub>oc</sub> Value (3)	Relative Losses		Toxicity		Soil Type (8)	Soil Leaching Rating (9)	Soil Runoff Rating (10)	Selected Pesticide (11)	Comments (12)
			Leaching RLPI (4)	Runoff RRPI (5)	MCL/HAL HALEQ (6)	Aquatic Toxicity (7)					

If the K<sub>oc</sub> value is 100 or less or if the RLPI value is 10 or less and the soil leach rating is high, then the pesticide has a high potential for leaching and should be used with extreme caution. Alternative pesticides and reduced rates should be considered if possible. Apply pesticide during periods with low potential for rainfall if possible.